

Using AMNet's Patient Reported Outcomes Measures (PROMs) to Implement ASAM's Clinical Practice Guideline on Alcohol Withdrawal Management

ASAM Recommendation

Applying AMNet's PROMS

Engage alcohol use disorder (AUD) patients with **comprehensive treatment** during alcohol withdrawal management



Track progress with:

Brief Addiction Monitor (BAM)

- 17-item scale assessing substance use, risk and protective factors

Treatment Effectiveness Assessment (TEA)

- 4-item recovery scale

Assess patients for **other substance use** and be prepared to treat **other potential withdrawal syndromes**

It may be helpful to use a validated scale that addresses other substance use



Other substance use can be tracked with:

TAPS Tool: alcohol and drug use screen & brief assessment

BAM

- Identifies drug use that may complicate alcohol withdrawal

- Evaluates the stability of recovery

- Informs need for additional support or interventions

Opioid withdrawal scales:

SOWS and **COWS**

Assess for **mental health conditions**
Psychiatric disorders may increase the risk for complicated alcohol withdrawal



PHQ-2+1: screens for depression and suicidal ideation

- Identifies need for diagnostic assessment

Evaluate **suicide risk** as part of the initial patient assessment



- Patients with suicidal ideation could be further screened with the **C-SSRS**

- It can help identify patients needing more urgent psychiatric evaluation