

<b>eMeasure Title</b>	<b>Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents</b>		
<b>eMeasure Identifier (Measure Authoring Tool)</b>	155	<b>eMeasure Version number</b>	5.0.000
<b>NQF Number</b>	0024	<b>GUID</b>	0b63f730-25d6-4248-b11f-8c09c66a04eb
<b>Measurement Period</b>	January 1, 20XX through December 31, 20XX		
<b>Measure Steward</b>	National Committee for Quality Assurance		
<b>Measure Developer</b>	National Committee for Quality Assurance		
<b>Endorsed By</b>	National Quality Forum		
<b>Description</b>	<p>Percentage of patients 3-17 years of age who had an outpatient visit with a Primary Care Physician (PCP) or Obstetrician/Gynecologist (OB/GYN) and who had evidence of the following during the measurement period. Three rates are reported.</p> <ul style="list-style-type: none"> <li>- Percentage of patients with height, weight, and body mass index (BMI) percentile documentation</li> <li>- Percentage of patients with counseling for nutrition</li> <li>- Percentage of patients with counseling for physical activity</li> </ul>		
<b>Copyright</b>	<p>Physician Performance Measure (Measures) and related data specifications were developed by the National Committee for Quality Assurance (NCQA).</p> <p>The Measures are copyrighted but can be reproduced and distributed, without modification, for noncommercial purposes (eg, use by healthcare providers in connection with their practices). Commercial use is defined as the sale, licensing, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial use of the Measures requires a license agreement between the user and NCQA. NCQA is not responsible for any use of the Measures.</p> <p>(C) 2008-2016 National Committee for Quality Assurance. All Rights Reserved.</p> <p>Limited proprietary coding is contained in the Measure specifications for user convenience. Users of proprietary code sets should obtain all necessary licenses from the owners of the code sets. NCQA disclaims all liability for use or accuracy of any CPT or other codes contained in the specifications.</p> <p>CPT(R) contained in the Measure specifications is copyright 2004-2015 American Medical Association. LOINC(R) copyright 2004-2015 Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms(R) (SNOMED CT[R]) copyright 2004-2015 International Health Terminology Standards Development Organisation. ICD-10 copyright 2015 World Health Organization. All Rights Reserved.</p>		
<b>Disclaimer</b>	<p>These performance Measures are not clinical guidelines and do not establish a standard of medical care, and have not been tested for all potential applications.</p> <p>THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.</p> <p>Due to technical limitations, registered trademarks are indicated by (R) or [R] and unregistered trademarks are indicated by (TM) or [TM].</p>		
<b>Measure Scoring</b>	Proportion		
<b>Measure Type</b>	Process		
<b>Stratification</b>	Report a total score, and each of the following strata: Stratum 1 - Patients age 3-11 Stratum 2 - Patients age 12-17		
<b>Risk Adjustment</b>	None		
<b>Rate Aggregation</b>	None		
<b>Rationale</b>	<p>One of the most important developments in pediatrics in the past two decades has been the emergence of a new chronic disease: obesity in childhood and adolescence. The rapidly increasing prevalence of obesity among children is one of the most challenging dilemmas currently facing pediatricians. National Health and Nutrition Examination Survey (NHANES) data from Cycle II (1976-1980) compared with data from Cycle III (1988-1994) documents an increase in the prevalence of obesity in all age, ethnic, and gender groups. NHANES data collected from 1999-2000 revealed a continued increase in the number of obese children. In that data collection, the prevalence of obesity (body mass index (BMI) &gt; 95th percentile) was 10 percent among children 2-5 years of age and 15 percent among children 6-19 years of age. When children at risk for obesity (BMI of 85th-94th percentile) were included, the prevalence increased to 20 percent and 30 percent, respectively. Therefore, &gt;1 of every 4 patients examined by pediatricians either is obese or is considered to be at high risk for developing this challenging health problem (O'Brien et al. 2004).</p> <p>In addition to the growing prevalence of obesity in children and adolescents, the number of overweight children at risk of becoming obese is also of great concern. Evidence suggests that overweight children and adolescents are more</p>		

	likely to become obese as adults. For example, one study found that approximately 80 percent of children who were overweight at age 10-15 years were obese adults at age 25 years (Whitaker et al. 1997). Another study found that 25 percent of obese adults were overweight as children. The latter study also found that if overweight begins before 8 years of age, obesity in adulthood is likely to be more severe (Freedman et al. 2001).
<b>Clinical Recommendation Statement</b>	<p>U.S. Preventive Services Task Force (2005) - Evidence is insufficient to recommend for or against routine screening for overweight in children and adolescents as a means to prevent adverse health outcomes (I rating).</p> <p>American Academy of Pediatrics (2004) - BMI should be calculated from the height and weight, and the BMI percentile should be calculated.</p> <p>American Medical Association (AMA), Centers for Disease Control and Prevention (CDC), Health Resources and Services Administration (HRSA) (2007) - At minimum, a yearly assessment of weight status in all children.</p> <p>Include calculation of height, weight (measured appropriately), and body mass index (BMI) for age and plotting of those measures on standard growth charts.</p> <p>The AAP and the American College of Clinical Endocrinology (ACCE) (Dorsey 2005) - Screen children for obesity using BMI and examine overweight children for obesity-related diseases.</p> <p>CDC (Baker 2005) - Using the percentile BMI for age and sex as the most appropriate and easily available method to screen for childhood overweight or at risk for overweight.</p> <p>Bright Futures (AAP) (Hagan 2008) - Calculate BMI at every visit.</p>
<b>Improvement Notation</b>	Higher score indicates better quality
<b>Reference</b>	O'Brien, S.H., R. Holubkov, E.C Reis. 2004. "Identification, evaluation, and management of obesity in an academic primary care center." <i>Pediatrics</i> 11:154-159.
<b>Reference</b>	Whitaker, R.C., J.A. Wright, M.S. Pepe, K.D. Seidel, W.H. Dietz. 1997. "Predicting obesity in young adulthood from childhood and parental obesity." <i>N Engl J Med</i> 37(13):869-873.
<b>Reference</b>	Freedman, D.S., L.K. Khan, W.H. Dietz, S.R. Srinivasan, G.S. Berenson. 2001. "Relationship of childhood overweight to coronary heart disease risk factors in adulthood: The Bogalusa Heart Study." <i>Pediatrics</i> 108:712-718.
<b>Reference</b>	U.S. Preventive Services Task Force (USPSTF). 2005. Screening and interventions for overweight in children and adolescents: recommendation statement. Rockville: Agency for Healthcare Research and Quality (AHRQ).
<b>Reference</b>	[AAP] National High Blood Pressure Education Program Working Group on High Blood Pressure in Children. 2004. "The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents." <i>Pediatrics</i> 114(2 Suppl):555-76.
<b>Reference</b>	AMA/HRSA/CDC Expert Committee on the Assessment, Prevention and Treatment of Child and Adolescent Overweight and Obesity. 2007. Recommendations on the assessment, prevention and treatment of child and adolescent overweight and obesity. Chicago: AMA.
<b>Reference</b>	Dorsey, K.B., C. Wells, H.M. Krumholz, J.C. Concato. 2005. "Diagnosis, evaluation, and treatment of childhood obesity in pediatric practice." <i>Arch Pediatr Adolesc Med</i> 159:632-638.
<b>Reference</b>	Baker, S., S. Barlow, W. Cochran, G. Fuchs, W. Klish, N. Krebs, R. Strauss, A. Tershakovec, J. Udall. 2005. "Overweight children and adolescents: a clinical report of the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition." <i>J Pediatr Gastroenterol Nutr</i> 40(5):533-43.
<b>Reference</b>	Hagan, J.F., Shaw J.S., Duncan P.M. eds. 2008. <i>Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, Third Edition</i> . Elk Grove: American Academy of Pediatrics.
<b>Definition</b>	None
<b>Guidance</b>	The visit must be performed by a PCP or OB/GYN. Because BMI norms for youth vary with age and sex, this measure evaluates whether BMI percentile is assessed rather than an absolute BMI value.
<b>Transmission Format</b>	TBD
<b>Initial Population</b>	Patients 3-17 years of age with at least one outpatient visit with a primary care physician (PCP) or an obstetrician/gynecologist (OB/GYN) during the measurement period
<b>Denominator</b>	Equals Initial Population
<b>Denominator Exclusions</b>	Patients who have a diagnosis of pregnancy during the measurement period
<b>Numerator</b>	Numerator 1: Patients who had a height, weight and body mass index (BMI) percentile recorded during the measurement period Numerator 2: Patients who had counseling for nutrition during a visit that occurs during the measurement period Numerator 3: Patients who had counseling for physical activity during a visit that occurs during the measurement period
<b>Numerator Exclusions</b>	Not Applicable

<b>Denominator Exceptions</b>	None
<b>Supplemental Data Elements</b>	For every patient evaluated by this measure also identify payer, race, ethnicity and sex

## Table of Contents

- [Population Criteria](#)
- [Data Criteria \(ODM Variables\)](#)
- [Data Criteria \(ODM Data Elements\)](#)
- [Supplemental Data Elements](#)
- [Risk Adjustment Variables](#)

## Population Criteria

### ----- Population Criteria 1 -----

- **Initial Population =**
  - AND: Age >= 3 year(s) at: "Measurement Period"
  - AND: Age < 17 year(s) at: "Measurement Period"
  - AND: \$OutpatientVisits
- **Denominator =**
  - AND: Initial Population
- **Denominator Exclusions =**
  - OR: "Diagnosis: Pregnancy" overlaps "Measurement Period"
- **Numerator =**
  - AND: "Physical Exam, Performed: BMI percentile (result)" during "Measurement Period"
  - AND: "Physical Exam, Performed: Height (result)" during "Measurement Period"
  - AND: "Physical Exam, Performed: Weight (result)" during "Measurement Period"
- **Numerator Exclusions =**
  - None
- **Denominator Exceptions =**
  - None
- **Stratifications =**
  - **Stratification 1 =**
    - AND: Age >= 3 year(s) at: "Measurement Period"
    - AND: Age < 11 year(s) at: "Measurement Period"
  - **Stratification 2 =**
    - AND: Age >= 11 year(s) at: "Measurement Period"
    - AND: Age < 17 year(s) at: "Measurement Period"

### ----- Population Criteria 2 -----

- **Initial Population =**
  - AND: Age >= 3 year(s) at: "Measurement Period"
  - AND: Age < 17 year(s) at: "Measurement Period"
  - AND: \$OutpatientVisits
- **Denominator =**
  - AND: Initial Population
- **Denominator Exclusions =**
  - OR: "Diagnosis: Pregnancy" overlaps "Measurement Period"
- **Numerator =**
  - AND: "Intervention, Performed: Counseling for Nutrition" during \$OutpatientVisits
- **Numerator Exclusions =**
  - None
- **Denominator Exceptions =**
  - None
- **Stratifications =**
  - **Stratification 1 =**
    - AND: Age >= 3 year(s) at: "Measurement Period"
    - AND: Age < 11 year(s) at: "Measurement Period"
  - **Stratification 2 =**
    - AND: Age >= 11 year(s) at: "Measurement Period"
    - AND: Age < 17 year(s) at: "Measurement Period"

### ----- Population Criteria 3 -----

- **Initial Population =**
  - AND: Age >= 3 year(s) at: "Measurement Period"
  - AND: Age < 17 year(s) at: "Measurement Period"
  - AND: \$OutpatientVisits
- **Denominator =**

- AND: Initial Population
- **Denominator Exclusions =**
  - OR: "Diagnosis: Pregnancy" overlaps "Measurement Period"
- **Numerator =**
  - AND: "Intervention, Performed: Counseling for Physical Activity" during \$OutpatientVisits
- **Numerator Exclusions =**
  - None
- **Denominator Exceptions =**
  - None
- **Stratifications =**
  - **Stratification 1 =**
    - AND: Age >= 3 year(s) at: "Measurement Period"
    - AND: Age < 11 year(s) at: "Measurement Period"
  - **Stratification 2 =**
    - AND: Age >= 11 year(s) at: "Measurement Period"
    - AND: Age < 17 year(s) at: "Measurement Period"

### Data Criteria (QDM Variables)

- **\$OutpatientVisits =**
  - Union of:
    - "Encounter, Performed: Face-to-Face Interaction"
    - "Encounter, Performed: Office Visit"
    - "Encounter, Performed: Preventive Care Services-Individual Counseling"
    - "Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17"
    - "Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17"
    - "Encounter, Performed: Preventive Care Services - Group Counseling"
    - "Encounter, Performed: Home Healthcare Services"
    - during "Measurement Period"

### Data Criteria (QDM Data Elements)

- "Diagnosis: Pregnancy" using "Pregnancy Grouping Value Set (2.16.840.1.113883.3.526.3.378)"
- "Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"
- "Encounter, Performed: Home Healthcare Services" using "Home Healthcare Services Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1016)"
- "Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"
- "Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17" using "Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)"
- "Encounter, Performed: Preventive Care Services - Group Counseling" using "Preventive Care Services - Group Counseling Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1027)"
- "Encounter, Performed: Preventive Care Services-Individual Counseling" using "Preventive Care Services-Individual Counseling Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1026)"
- "Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17" using "Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)"
- "Intervention, Performed: Counseling for Nutrition" using "Counseling for Nutrition Grouping Value Set (2.16.840.1.113883.3.464.1003.195.12.1003)"
- "Intervention, Performed: Counseling for Physical Activity" using "Counseling for Physical Activity Grouping Value Set (2.16.840.1.113883.3.464.1003.118.12.1035)"
- "Physical Exam, Performed: BMI percentile" using "BMI percentile Grouping Value Set (2.16.840.1.113883.3.464.1003.121.12.1012)"
- "Physical Exam, Performed: Height" using "Height Grouping Value Set (2.16.840.1.113883.3.464.1003.121.12.1014)"
- "Physical Exam, Performed: Weight" using "Weight Grouping Value Set (2.16.840.1.113883.3.464.1003.121.12.1015)"

### Supplemental Data Elements

- "Patient Characteristic Ethnicity: Ethnicity" using "Ethnicity CDCREC Value Set (2.16.840.1.114222.4.11.837)"
- "Patient Characteristic Payer: Payer" using "Payer SOP Value Set (2.16.840.1.114222.4.11.3591)"
- "Patient Characteristic Race: Race" using "Race CDCREC Value Set (2.16.840.1.114222.4.11.836)"
- "Patient Characteristic Sex: ONC Administrative Sex" using "ONC Administrative Sex AdministrativeGender Value Set (2.16.840.1.113762.1.4.1)"

### Risk Adjustment Variables

- None

Measure Set	None
-------------	------