

This table is dedicated to Mental Health Non-Psychiatrist focused performance measures. These measures examine the treatment or evaluation of psychiatric symptoms and disorders by health professionals other than psychiatrists. When patients with a mental health condition such as Major Depression or ADHD are measured in the primary care setting for example recommended guidelines are not always followed. A current area of focus in psychiatric performance measurement is patient outcome indicators as measured by the PHQ-9 measurement tool. This tool can be utilized in the primary care setting and mental health care setting. More detailed information about each measure maybe found on the "All Measures" Spreadsheet.

Origin	Title	Description
NQF #0106	Diagnosis of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients newly diagnosed with attention deficit hyperactivity disorder (ADHD) whose medical record contains documentation of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) or Diagnostic and Statistical Manual for Primary Care (DSM-PC) criteria being addressed.
NQF #0107	Management of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients treated with psycho-stimulant medication for the diagnosis of attention deficit hyperactivity disorder (ADHD) whose medical record contains documentation of a follow-up visit at least twice a year.
NQMC	Depression : percent of patients with a diagnosis of depression with documented self-management goals set within the last 12 months.	This population-based measure is used in primary care settings to assess the percent of patients with a diagnosis of depression with documented self-management goals set within the last 12 months. Patient education and self-management skills have been important components of collaborative model interventions (Katon W, et al., JAMA 1995; Olivarius N, et al., BMJ. 2001; Renders CM, et al., Diabetes Care. 2001; Norris SL, et al., Diabetes Care. 2001; Maly RM, et al., Jrnl Amer Ger Soc 2002; Heisler M, et al., J Gen Inter Med 2002; Lorig KR, et al., Medical Care 1999).
NQMC	Depression : the percentage of patients on the diabetes register and/or coronary heart disease (CHD) register for whom case finding for depression has been undertaken on one occasion during the previous 15 months using two standard screening questions.	This measure is used to assess the percentage of patients on the diabetes register and/or coronary heart disease (CHD) register for whom case finding for depression has been undertaken on one occasion during the previous 15 months using two standard screening questions.
NQMC	Major depression in adults in primary care : percentage of patients who have a depression follow-up contact within three months of initiating treatment.	This measure is used to assess the percentage of patients who have a depression follow-up contact* within three months of initiating treatment. *Contact = an office visit or phone contact with physician or other care provider.
NQMC	Major depression in adults in primary care : percentage of patients with a new diagnosis of major depression, with documentation of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV TR) criteria within the three months prior to initial diagnosis.	This measure is used to assess the percentage of patients with a new diagnosis of major depression, with documentation of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV TR) criteria within the three months prior to initial diagnosis.
NQMC	Major depression in adults in primary care : percentage of patients who commit suicide at any time while under depression management with a primary care physician.	This measure is used to assess the percentage of patients who commit suicide at any time while under depression management with a primary care physician.

NQMC	Major depression in adults in primary care : percentage of patients with diabetes with documentation of screening for depression.	This measure is used to assess the percentage of patients with diabetes with documentation in the medical record of screening for depression.
NQF #1401	Maternal Depression Screening	The percentage of children who turned 6 months of age during the measurement year who had documentation of a maternal depression screening for the mother.
CQAIMH	Substance Abuse Education in Primary Care	