

This table is dedicated to performance measures around substance use disorder or dual diagnosis. These measures assess care focusing on *retention rates* (assessing the amount of time committed to treatment against length of time until relapse), *screening*, and *treatment management* among others. More detailed information about each measure maybe found on the "All Measures" Spreadsheet.

Origin	Title	Description
CQAIMH	<a href="#">Continuity of Care for Dual Diagnoses</a>	
NQF #0004	<a href="#">Initiation and Engagement of Alcohol and Other Drug Dependence Treatment</a>	The percentage of adolescent and adult members with a new episode of alcohol or other drug (AOD) dependence who received the following.  a. Initiation of AOD Treatment. The percentage of members who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis. b. Engagement of AOD Treatment. The percentage of members who initiated treatment and who had two or more additional services with a diagnosis of AOD within 30 days of the initiation visit. <b>[Please note this measure is appropriate for Health Plan or State Level Measurement only.]</b>
CQAIMH	<a href="#">Substance Abuse Education in Primary Care</a>	
CQAIMH	<a href="#">1-Month Retention Rate for Chemical Dependency Treatment</a>	
CQAIMH	<a href="#">1-Year Retention Rate for Chemical Dependency Treatment</a>	
CQAIMH	<a href="#">3-Month Retention Rate for Chemical Dependency Treatment</a>	
CQAIMH	<a href="#">60-day Continuation of Substance Abuse Treatment</a>	
CQAIMH	<a href="#">6-Month Retention Rate for Chemical Dependency Treatment</a>	
CQAIMH	<a href="#">Continuation after Substance-Related Treatment Initiation</a>	
CQAIMH	<a href="#">Maintenance Pharmacotherapy for Substance Abuse</a>	
CQAIMH	<a href="#">Multiple Outpatient Visits after Substance-Related Hospitalization</a>	
CQAIMH	<a href="#">Outpatient Follow-up after Initial Substance-Related Visit (2 or more visits)</a>	
CQAIMH	<a href="#">Program Completion for Chemical Dependency Treatment</a>	
CQAIMH	<a href="#">Substance Abuse Maintenance Treatment</a>	
NQMC	<a href="#">Substance use disorders</a> : percentage of patients aged 18 years and older with a diagnosis of current substance abuse or dependence who were screened for depression within the 12 month reporting period.	This measure is used to assess the percentage of patients aged 18 years and older with a diagnosis of current substance abuse or dependence who were screened for depression within the 12 month reporting period.
NQMC	<a href="#">Substance use disorders</a> : percentage of patients aged 18 years and older with a diagnosis of current alcohol dependence who were counseled regarding psychosocial AND pharmacologic treatment options for alcohol dependence within the 12 month reporting period.	This measure is used to assess the percentage of patients aged 18 years and older with a diagnosis of current alcohol dependence who were counseled regarding psychosocial AND pharmacologic treatment options for alcohol dependence within the 12 month reporting period.

<b>NQMC</b>	<a href="#">Assessment of smoking, substance abuse, safety, and firearms risks in the family</a> : average percentage of recommended topics assessed.	This measure is used to assess the average percentage of recommended topics the health care providers asked about related to smoking, substance abuse, safety, and firearms in the home.
<b>NQMC</b>	<a href="#">Preventive screening and counseling on risky behaviors</a> : average proportion saying "yes" to ten items about whether provider(s) discussed/screened on smoking, alcohol use, helmet use, drunk driving, chewing tobacco, street drugs, steroid pills, sexual/physical abuse, violence, guns.	This measure assesses the average proportion of "yes" responses to ten items about whether provider(s) discussed/screened on smoking, alcohol use, helmet use, drunk driving, chewing tobacco, street drugs, steroid pills, sexual/physical abuse, violence, and guns among young adults.
<b>PCPI</b>	<a href="#">Substance Use Disorders Measure #2</a> : Counseling Regarding Psychosocial and Pharmacologic Treatment Options for Opioid Addiction	Percentage of patients aged 18 years and older with a diagnosis of current opioid addiction who were counseled regarding psychosocial AND pharmacologic treatment options for opioid addiction within the 12 month reporting period
<b>The Washington Circle Group</b>	<a href="#">Treatment Engagement</a>	Treatment engagement is an intermediate step between initially accessing care (in the first visit) and completing a full course of treatment. This measure is designed to assess the degree to which members engage in treatment with two additional AOD treatments within 30 days after initiating treatment.
<b>NQMC</b>	<a href="#">Substance use</a> : percent of patients who are identified with alcohol or drug disorder who receive a prescription for FDA-approved medication for alcohol or drug use disorder or a referral for addictions treatment.	This measure is used to assess the percent of hospitalized patients 18 years of age and older who are identified with an alcohol or drug use disorder who received a prescription at discharge for Food and Drug Administration (FDA)-approved medication for treatment of alcohol or drug use disorder OR a referral for addictions treatment.